

2007 Extension 1 Solution (by Terry Lee)

Q1

$$\begin{aligned} \text{(a)} \quad (1 + \sqrt{5})^3 &= 1 + 3\sqrt{5} + 3(\sqrt{5})^2 + (\sqrt{5})^3 \\ &= 1 + 3\sqrt{5} + 15 + 5\sqrt{5} \\ &= 16 + 8\sqrt{5}. \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad x &= \frac{3 \times 4 + 2 \times 19}{3 + 2} = 10 \\ y &= \frac{3 \times 5 + 2 \times (-5)}{3 + 2} = 1. \end{aligned}$$

$$\text{(c)} \quad \frac{d}{dx}(\tan^{-1}(x^4)) = \frac{4x^3}{1+x^8}.$$

(d) For $y = x^3 + 1$, $y' = 3x^2$. When $x = 1$, $m_1 = 3$.

$$\text{For } x - 2y + 3 = 0, m_2 = \frac{1}{2}.$$

$$\tan \alpha = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \left| \frac{3 - \frac{1}{2}}{1 + 3 \times \frac{1}{2}} \right| = \frac{5}{5} = 1, \therefore \alpha = \frac{\pi}{4}.$$

(e) Let $u = 25 - x^2$, $du = -2x dx$.

When $x = 3$, $u = 16$; When $x = 4$, $u = 9$.


$$\begin{aligned} \int_3^4 \frac{2x}{\sqrt{25-x^2}} dx &= \int_{16}^9 \frac{-du}{\sqrt{u}} \\ &= 2 \left[\sqrt{u} \right]_9^{16} \\ &= 2(4 - 3) \\ &= 2. \end{aligned}$$

Q2

$$\begin{aligned} \text{(a) LHS} &= \frac{1 - \frac{1-t^2}{1+t^2}}{\frac{2t}{1+t^2}} \\ &= \frac{1+t^2 - 1+t^2}{2t} \\ &= \frac{2t^2}{2t} \\ &= t = \text{RHS}. \end{aligned}$$

(b) (i) See graph.

(ii) $0 \leq y \leq 2\pi$.



$$\text{(c) } P(2) = 0 \text{ gives } 4 + 2a + b = 0. \quad (1)$$

$$P(-1) = 18 \text{ gives } 18 = 1 - a + b, \therefore 17 + a - b = 0 \quad (2)$$

$$(1) + (2) \text{ gives } 21 + 3a = 0, \therefore a = -7.$$

Substituting $a = -7$ to (1) gives $b = 10$.

$$\text{(d) (i) } a = \frac{dv}{dt} = 50 \times 0.2e^{-0.2t}.$$

When $t = 10$, $a = 10e^{-2} = 1.4 \text{ m/s}^2$.

$$\text{(ii) } \frac{dx}{dt} = 50(1 - e^{-0.2t})$$

$$x = \int_0^{10} 50(1 - e^{-0.2t}) dt$$

$$= 50 \left[t + \frac{e^{-0.2t}}{0.2} \right]_0^{10}$$


$$= 50 \left[\left(10 + \frac{e^{-2}}{0.2} \right) - \frac{1}{0.2} \right]$$

$$= 284 \text{ m}.$$

Q3

$$\begin{aligned}
 \text{(a) } V &= \int_0^3 \pi y^2 dx \\
 &= \pi \int_0^3 \frac{1}{9+x^2} dx \\
 &= \pi \left[\frac{1}{3} \tan^{-1} \frac{x}{3} \right]_0^3 \\
 &= \frac{\pi}{3} \times \frac{\pi}{4} \\
 &= \frac{\pi^2}{12}.
 \end{aligned}$$

- (b) (i) Vertical asymptote $x = 4$.
Horizontal asymptote $y = 1$.



Solving $\frac{x-2}{x-4} = 3$ gives $x-2 = 3x-12$

$$2x = 10$$

$$x = 5.$$

$$\therefore \frac{x-2}{x-4} \leq 3 \text{ when } x < 4 \text{ or } x \geq 5.$$

(c) (i) $\ddot{x} = \frac{d}{dx} \left(\frac{1}{2} v^2 \right), \therefore v^2 = 2 \int \ddot{x} dx$

$$v^2 = 2 \int -e^{-2x} dx = e^{-2x} + C.$$

When $t = 0, x = 0, v = 1, \therefore 1 = 1 + C, \therefore C = 0.$

$$\therefore v^2 = e^{-2x}.$$

$$\therefore v = \pm e^{-x}.$$

Initially, $v > 0, \therefore v = e^{-x}.$

(ii) $\frac{dx}{dt} = e^{-x}$

$$\frac{dt}{dx} = e^x.$$

$$t = e^x + C.$$

$t = 0, x = 0, \therefore C = -1.$

$$\therefore t = e^x - 1.$$

$$e^x = t + 1.$$

$$x = \ln(t + 1).$$

Q4

(a) (i) $(0.1)^2 = 0.01$

(ii) ${}^{20}C_2 (0.1)^2 (0.9)^{18} = 0.285$

(iii) $1 - \Pr(x = 0, 1, 2)$

$$\begin{aligned}
 &= 1 - 0.9^{20} - {}^{20}C_1 (0.1)(0.9)^{19} - {}^{20}C_2 (0.1)^2 (0.9)^{18} \\
 &= 0.32.
 \end{aligned}$$

- (b) Let $n = 1, 7 + 5 = 12$, which is divisible by 12,

\therefore The statement is true for $n = 1$.

Assume the statement is true for $n = k$,

i.e. $7^{2k-1} + 5 = 12M$, where M is an integer.

$$\therefore 7^{2n-1} = 12M - 5$$

$$\therefore 7^{2n+1} + 5 = 49(7^{2n-1}) + 5$$

$$= 49(12M - 5) + 5$$

$$= 588M - 240$$

$$= 12(49M - 20), \text{ which is a multiple of 12.}$$

\therefore The statement is true for all $n = k + 1$.

\therefore The statement is true for all $n \geq 1$.

- (c) (i) $\angle QXB = \angle DXP$ (vertically opposite)

$$\angle DXP = \angle XAP \text{ (both } = 90^\circ - \angle XDP, \text{ angle sum in } \Delta)$$

$$\angle XAP = \angle QBX \text{ (angles subtending the same arc are equal)}$$

$$\therefore \angle QXB = \angle QBX.$$

(ii) From (i), ΔBXQ is isosceles, $\therefore QB = QX$ (base angles in isosceles Δ).

Similarly, $\angle QXC = \angle QCX, \therefore \Delta CXQ$ is isosceles,

$$\therefore QC = QX.$$

$$\therefore QB = QC.$$

$$\therefore Q \text{ bisects } BC.$$

Q5

$$\begin{aligned} \text{(a) (i) Area } (\triangle OPT) &= \frac{1}{2} OP \cdot TP \\ &= \frac{1}{2} r^2 \tan \theta, \text{ since } TP = r \tan \theta. \end{aligned}$$

$$\text{Area (sector } OPQ) = \frac{1}{2} r^2 \theta.$$

$$\text{Area } (\triangle OPT) = 2 \times \text{Area (sector } OPQ),$$

$$\therefore \frac{1}{2} r^2 \tan \theta = 2 \times \frac{1}{2} r^2 \theta.$$

$$\therefore \tan \theta = 2\theta.$$

$$\text{(ii) Let } f(x) = 2\theta - \tan \theta.$$

$$f'(x) = 2 - \sec^2 \theta.$$

$$\text{By Newton's method, } x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$\begin{aligned} x_1 &= 1.15 - \frac{2 \times 1.15 - \tan 1.15}{2 - \sec^2 1.15} \\ &= 1.1664. \end{aligned}$$

- (b) Group the four children together, there are 3 groups,
 $\therefore 3!4!$ ways to arrange so that the children stay together.
 $\therefore \text{Pr}(\text{the 4 children stay together}) = \frac{3!4!}{6!} = \frac{1}{5}.$

- (c) Let $u = x, v = y,$

$$\sin^{-1} x + \frac{1}{2} \cos^{-1} y = \frac{\pi}{3} \quad (1)$$

$$3 \sin^{-1} x - \frac{1}{2} \cos^{-1} y = \frac{2\pi}{3} \quad (2)$$

$$(1) + (2) \text{ gives } 4 \sin^{-1} x = \pi.$$

$$\therefore \sin^{-1} x = \frac{\pi}{4}, \therefore x = \frac{\sqrt{2}}{2}.$$

$$\text{Substituting } \sin^{-1} x = \frac{\pi}{4} \text{ into (1),}$$

$$\frac{1}{2} \cos^{-1} y = \frac{\pi}{3} - \frac{\pi}{4}$$

$$\therefore \cos^{-1} y = \frac{\pi}{6}, \therefore y = \frac{\sqrt{3}}{2}.$$

- (d) (i) Substituting $(2aq, aq^2)$ into the equation of $PQ,$

$$2aq + p(aq^2) - 2ap - ap^3 = 0$$

$$2q + pq^2 - 2p - p^3 = 0$$

$$2(p - q) + p(q^2 - p^2) = 0$$

$$2(p - q) + p(p - q)(p + q) = 0$$

$$2 + p(p + q) = 0, \text{ since } p \neq q.$$

$$2 + p^2 + pq = 0.$$

$$\text{(ii) If } OP \perp OQ \text{ then } \frac{ap^2}{2ap} \times \frac{aq^2}{2aq} = -1, \therefore pq = -4.$$

$$\text{Substituting to part (i), } 2 + p^2 - 4 = 0.$$

$$\therefore p^2 = 2.$$

Q6

$$\text{(a) (i) } x = 3 + \sqrt{3} \sin 2t - \cos 2t$$

$$\dot{x} = 2\sqrt{3} \cos 2t + 2 \sin 2t$$

$$\ddot{x} = -4\sqrt{3} \sin 2t + 4 \cos 2t$$

$$= -4(\sqrt{3} \sin 2t - \cos 2t)$$

$$= -4(x - 3).$$

$$\text{(ii) Period} = \frac{2\pi}{n} = \frac{2\pi}{2} = \pi \text{ s.}$$

$$\text{(iii) } \dot{x} = 2\sqrt{3} \cos 2t + 2 \sin 2t$$

$$= 4 \left(\frac{\sqrt{3}}{2} \cos 2t + \frac{1}{2} \sin 2t \right)$$

$$= 4 \left(\cos 2t \cos \frac{\pi}{6} + \sin 2t \sin \frac{\pi}{6} \right)$$

$$= 4 \cos \left(2t - \frac{\pi}{6} \right).$$

$$\text{(iv) When } \dot{x} = \pm 2,$$

$$\pm 2 = 4 \cos \left(2t - \frac{\pi}{6} \right)$$

$$\cos \left(2t - \frac{\pi}{6} \right) = \pm \frac{1}{2}$$

$$2t - \frac{\pi}{6} = \pm \frac{\pi}{3} + k2\pi, \pm \frac{2\pi}{3} + k2\pi.$$

$$2t = \pm \frac{\pi}{3} + \frac{\pi}{6} + k2\pi, \pm \frac{2\pi}{3} + \frac{\pi}{6} + k2\pi.$$

$$t = \pm \frac{\pi}{6} + \frac{\pi}{12} + k\pi, \pm \frac{\pi}{3} + \frac{\pi}{12} + k\pi.$$

$$\text{For } 0 \leq t \leq \pi, t = \frac{\pi}{4}, \frac{5\pi}{12}, \frac{3\pi}{4}, \frac{11\pi}{12}.$$

- (b) (i) $f'(x) = e^x + e^{-x} > 0$ always, $\therefore f(x)$ is increasing.

$$\text{(ii) } f^{-1} : x = e^y - e^{-y} \quad (1)$$

$$x = e^y - \frac{1}{e^y}.$$

$$xe^y = e^{2y} - 1.$$

$$e^{2y} - xe^y - 1 = 0$$

$$e^y = \frac{x \pm \sqrt{x^2 + 4}}{2}.$$

$$\text{Since } e^y > 0 \text{ for all real } x, e^y = \frac{x + \sqrt{x^2 + 4}}{2}.$$

$$\therefore y = \ln \frac{x + \sqrt{x^2 + 4}}{2}. \quad (2)$$

- (iii) Substituting $x = 5$ in (1) gives the same answer as substituting $x = 5$ in (2).

$$\therefore \text{Solving } e^x - e^{-x} = 5 \text{ gives } x = \ln \frac{5 + \sqrt{29}}{2} = 1.65$$

Q7

(a) (i) $y = kx^n, \therefore y' = nkx^{n-1}$.

When $x = a, m_1 = nka^{n-1}$.

$$y = \ln x, y' = \frac{1}{x}$$

When $x = a, m_2 = \frac{1}{a}$.

If these two gradients are equal, $nka^{n-1} = \frac{1}{a}$.

$$\therefore nka^n = 1.$$

$$\therefore a^n = \frac{1}{nk} \quad (1)$$

(ii) When $x = a, ka^n = \ln a$.

$$k \times \frac{1}{nk} = \ln a, \text{ on substituting } a^n = \frac{1}{nk}$$

$$\ln a = \frac{1}{n}$$

$$a = e^{\frac{1}{n}}$$

$$a^n = e.$$

$$\therefore (1) \text{ becomes } e = \frac{1}{nk}, \therefore k = \frac{1}{ne}.$$

(b) (i) $t = \frac{x}{14 \cos \theta}$.

$$\begin{aligned}
 y &= 14 \left(\frac{x}{14 \cos \theta} \right) \sin \theta - 4.9 \left(\frac{x}{14 \cos \theta} \right)^2 \\
 &= x \tan \theta - \frac{1}{40} x^2 \sec^2 \theta \\
 &= x \tan \theta - \frac{1}{40} x^2 (1 + \tan^2 \theta) \\
 &= mx - \left(\frac{1+m^2}{40} \right) x^2, \text{ by letting } \tan \theta = m.
 \end{aligned}$$

(ii) When $x = 10, y = h$,

$$h = 10m - \left(\frac{1+m^2}{40} \right) 100$$

$$h = 10m - 2.5(1+m^2).$$

$$2.5m^2 - 10m + 2.5 + h = 0.$$

$$m = \frac{10 \pm \sqrt{100 - 10(2.5 + h)}}{5}$$

$$= \frac{10 \pm \sqrt{75 - 10h}}{5}$$

$$= \frac{10 \pm 5\sqrt{3 - 0.4h}}{5}$$

$$= 2 \pm \sqrt{3 - 0.4h}.$$

Since $3 - 0.4h \geq 0, h \leq \frac{3}{0.4} = 7.5$ m.

(iii) Given $m = 2 \pm \sqrt{3 - 0.4h}$.

When $h = 5.9, m = 2 \pm \sqrt{3 - 0.4 \times 5.9} = 2.8$ or 1.2 .

When $h = 3.9, m = 2 \pm \sqrt{3 - 0.4 \times 3.9} = 0.8$ or 3.2 .

\therefore One interval is $2.8 \leq m \leq 3.2$, and the other interval is $0.8 \leq m \leq 1.2$.

(iv) When $y = 0, mx - \left(\frac{1+m^2}{40} \right) x^2 = 0$

$$x \left(m - \frac{1+m^2}{40} x \right) = 0$$

$$x = \frac{40m}{1+m^2} \text{ m.}$$

When $m = 2.8, x = 12.7$ m

When $m = 3.2, x = 11.4$ m

The width of this interval is $12.7 - 11.4 = 1.3$ m.

When $m = 1.2, x = 19.7$ m

When $m = 0.8, x = 19.5$ m

But m is the gradient of the angle of projection, so when $0.8 \leq m \leq 1.2$, the maximum range is obtained when $\theta = 45^\circ$ (i.e. $m = 1$), \therefore When $m = 1, x = 20$ m.

The width of this interval is $20 - 19.5 = 0.5$ m.